



# CHILD ADMINISTRATION INSTRUCTIONS



1 Put person on their side where they can't fall.



2 Get medicine.



3 Get syringe. *Note: seal pin is attached to the cap.*



4 Push up with thumb and pull to remove cap from syringe. **Be sure seal pin is removed with the cap.**



5 Lubricate rectal tip with lubricating jelly.




6 Turn person on side facing you.



7 Bend upper leg forward to expose rectum.




8 Separate buttocks to expose rectum.




9 Gently insert syringe tip into rectum. *Note: rim should be snug against rectal opening.*

**SLOWLY...**


**COUNT OUT LOUD TO THREE...1...2...3**



10 Slowly count to 3 while gently pushing plunger in until it stops.




11 Slowly count to 3 before removing syringe from rectum.



12 Slowly count to 3 while holding buttocks together to prevent leakage.

**ONCE DIASTAT® IS GIVEN**



13

Keep person on the side facing you, note time given, and continue to observe.

**CALL FOR HELP IF ANY OF THE FOLLOWING OCCUR**

- Seizure(s) continues 15 minutes after giving DIASTAT® or per the doctor's instructions:

- Seizure behavior is different from other episodes
- You are alarmed by the frequency or severity of the seizure(s)
- You are alarmed by the color or breathing of the person
- The person is having unusual or serious problems

Local emergency number: \_\_\_\_\_ Doctor's number: \_\_\_\_\_  
(Please be sure to note if your area has 911)

Information for emergency squad: Time DIASTAT® given: \_\_\_\_\_ Dose: \_\_\_\_\_

**DIASTAT® Indication**

DIASTAT® AcuDial™ (diazepam rectal gel) is a gel formulation of diazepam intended for rectal administration in the management of selected, refractory patients with epilepsy, on stable regimens of AEDs, who require intermittent use of diazepam to control bouts of increased seizure activity, for patients 2 years and older.

**Important Safety Information**

In clinical trials with DIASTAT®, the most frequent adverse event was somnolence (23%). Less frequent adverse events reported were dizziness, headache, pain, vasodilatation, diarrhea, ataxia, euphoria, incoordination, asthma, rash, abdominal pain, nervousness, and rhinitis (1%-5%).


D955-0308

**Diastat®** (diazepam rectal gel) **Diastat AcuDial™** (diazepam rectal gel) 

**DISPOSAL INSTRUCTIONS ON REVERSE SIDE**

## Procedure

- Cushion the child's head
- Loosen any tight neckwear
- Turn the child on his or her side
- Do not hold the child down or restrain the person
- Do not place anything in the mouth or try to pry the teeth apart. The child is not in danger of swallowing his or her tongue
- Observe seizure characteristics -- length, type of movements, direction of head or eye turning. These characteristics may help the doctor diagnose the type of seizure



## Call 911

- First time seizure If you are unaware of a seizure disorder: it is a first time seizure!
- Seizure over 5 minutes
- Administer Diastat only with training
- Call Parents...inform principal and supervisor